

Name of the Student: _____

Max. Marks : 17 Marks

Time : 17 Minutes

Q1.

An excess of a given reagent is added to each of the following pairs of aqueous metal ions.

For each metal ion, state the initial colour of the solution and the final observation that you would make.

In each case, write an overall equation for the formation of the final product from the initial aqueous metal ion.

- (a) An excess of aqueous sodium carbonate is added to separate aqueous solutions containing $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ and $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$.

(5)

- (b) An excess of concentrated hydrochloric acid is added to separate aqueous solutions containing $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ and $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$.

(4)

- (c) An excess of dilute aqueous sodium hydroxide is added to separate aqueous solutions containing $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ and $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$.

(4)

- (d) An excess of dilute aqueous ammonia is added to separate aqueous solutions containing $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ and $[\text{Ag}(\text{H}_2\text{O})_2]^+$

(4)

(Total 17 marks)