

Name of the Student: _____

Max. Marks : 16 Marks

Time : 16 Minutes

Q1.


A student carried out an experiment to find the temperature rise for a reaction between hydrochloric acid and sodium hydroxide solution.

- The student used a measuring cylinder to place 50 cm^3 of $0.400 \text{ mol dm}^{-3}$ hydrochloric acid into a glass beaker.
- The student recorded the temperature at one-minute intervals for three minutes.
- At the fourth minute the student added 50 cm^3 of $0.400 \text{ mol dm}^{-3}$ sodium hydroxide solution and stirred to mix the solutions, but did not record the temperature.
- The student recorded the temperature at one-minute intervals for a further eight minutes.

The results are shown in the table.

Time / min	0	1	2	3	4	5	6	7	8	9	10	11	12
Temperature / $^{\circ}\text{C}$	19.8	19.8	19.8	19.8		21.4	21.7	21.6	21.5	21.4	21.3	21.2	21.1

- (a) Plot a graph of temperature against time on the grid below.



Use your graph to find the temperature rise, ΔT , at the fourth minute.
Show your working on the graph by drawing suitable lines of best fit.

- (b) The uncertainty in each of the temperature readings from the thermometer used in this experiment was $\pm 0.1^\circ \text{C}$

Calculate the percentage uncertainty in the value for the temperature rise.

Percentage uncertainty _____

(1)

- (c) Suggest a change to the experiment that would minimise heat loss.

(1)

- (d) Suggest and explain another change to the experiment that would decrease the percentage uncertainty in the use of the same thermometer.

(2)

- (e) A second student completed an experiment to determine the enthalpy of neutralisation for the reaction between ethanedioic acid solution (HOOCCOOH) and potassium hydroxide solution.

The student added 25 cm^3 of 0.80 mol dm^{-3} ethanedioic acid solution to 75 cm^3 of 0.60 mol dm^{-3} potassium hydroxide solution.

The temperature increased by 3.2°C

Give an equation for the reaction between ethanedioic acid solution and potassium hydroxide solution.

Calculate the enthalpy change (ΔH) per mole of water formed in this reaction.

Assume that the specific heat capacity of the reaction mixture is $4.2 \text{ J K}^{-1} \text{ g}^{-1}$

Assume that the density of the reaction mixture is 1.00 g cm^{-3}

Equation _____

- (f) In a similar experiment to that in **part (e)**, the enthalpy of neutralisation for the reaction between sulfuric acid and potassium hydroxide solution was found to be $-57.0 \text{ kJ mol}^{-1}$ per mole of water formed.

Suggest an explanation for the difference between this value and your answer to **part (e)**.

(If you were unable to obtain an answer to **part (e)** you should assume a value of $-28.5 \text{ kJ mol}^{-1}$. This is **not** the correct answer.)

(2)

(Total 16 marks)