Practice Question Set For A-Level
Subject : Chemistry
Paper-1 Topic: Physical 1

Merit Minds
www.merit-minds.com
Exam Preparation and Free Resources

Name of the Student:_____
Max. Marks : 25 Marks **Time : 25 Minutes**

Mark Schemes

Q1.

(a) Particles are in maximum state of order

(or perfect order or completely ordered or perfect crystal or minimum disorder or no disorder)

(entropy is zero at 0 k by definition)

1

(b) (Ice) melts

(or freezes or changes from solid to liquid or from liquid to solid)

1

(c) Increase in disorder

1

Bigger (at T_2)

1

Second mark only given if first mark has been awarded

(d) (i) Moles of water = 1.53/18 (= 0.085)

1

Heat change per mole = 3.49/0.085 = 41.1 (kJ mol^{-1})

(allow 41 to 41.1, two sig. figs.)

(penalise –41 (negative value), also penalise wrong units but allow kJ only)

1

(ii) $\Delta G = \Delta H - T\Delta S$

1

(iii) $\Delta H = T\Delta S$ or $\Delta S = \Delta H/T$

(penalise if contradiction)

1

ΔS = 41.1/373 = 0.110 kJ K^{-1} (mol^{-1}) (or 110 (J K^{-1} (mol^{-1}))

(allow 2 sig. figs.)

(if use value given of 45, answer is 0.12 (or 120 to 121)

(if ΔH is negative in (d) (i), allow negative answer)

(if ΔH is negative in (d) (i), allow positive answer)

(if ΔH is positive in (d) (i), penalise negative answer)

1

Correct units as above (mol^{-1} not essential)

1

Downloaded from www.merit-minds.com

[10]

Q2.

(a) *M1* $K_p = (_PY)^3 . (_PZ)^2 / (_PW)^2 . (_PX)$ *NB [ ] wrong*

1

M2 temperature

1

M3 increase

1

M4 particles have more energy or greater velocity/speed

1

M5 more collisions with $E > E_a$ or more successful collisions

1

M6 Reaction exothermic or converse

1

M7 Equilibrium moves in the left

1

Marks for other answers

Increase in pressure or concentration	*allow M1, M5, M6*	*Max 3*
Addition of a catalyst;	*allow M1, M5, M6*	*Max 3*
Decrease in temperature;	*allow M1, M2, M6*	*Max 3*
Two or more changes made;	*allow M1, M6*	*Max 2*

(b) (i) Advantage; reaction goes to completion, not reversible or faster

1

Disadvantage; reaction vigorous/dangerous

(exothermic must be qualified)

or HCl(g) evolved/toxic
or CH_3COCl expensive

NB Allow converse answers
Do not allow reactions with other reagents e.g. water or ease of separation

1

(ii) $\Delta S = \Sigma S$ products $- \Sigma S$ reactants

1

$\Delta S = (259 + 187) - (201 + 161)$

1

$\Delta S = 84$ (JK^{-1} mol^{-1}) *(Ignore units)*

*Allow – 84 to score **(1)** mark*

1

$\Delta G = \Delta H - T\Delta S$

1

$= -21.6 - 298 \times 84/1000$
$= -46.6$ kJ mol^{-1} or $-46\,600$ J mol^{-1}

1

*Allow **(2)** for – 46.6 without units*

Downloaded from www.merit-minds.com

(Mark ΔG consequentially to incorrect ΔS)
(e.g. ΔS = –84 gives ΔG = +3.4 kJ mol^{-1})

1

[15]

Downloaded from www.merit-minds.com