

Name of the Student: _____

Max. Marks : 21 Marks

Time : 21 Minutes

Q1.

Acyl chlorides and acid anhydrides are important compounds in organic synthesis.


- (a) Outline a mechanism for the reaction of $\text{CH}_3\text{CH}_2\text{COCl}$ with CH_3OH and name the organic product formed.

Mechanism

Name of organic product _____

(5)

- (b) A polyester was produced by reacting a diol with a diacyl chloride. The repeating unit of the polymer is shown below.



- (i) Name the diol used.

(1)

- (ii) Draw the displayed formula of the diacyl chloride used.

(1)

- (iii) A shirt was made from this polyester. A student wearing the shirt accidentally splashed aqueous sodium hydroxide on a sleeve. Holes later appeared in the sleeve where the sodium hydroxide had been.


Name the type of reaction that occurred between the polyester and the aqueous sodium hydroxide. Explain why the aqueous sodium hydroxide reacted with the polyester.

Type of reaction _____

Explanation _____

(3)

- (c) (i) Complete the following equation for the preparation of aspirin using ethanoic anhydride by writing the structural formula of the missing product.



(1)

- (ii) Suggest a name for the mechanism for the reaction in part (c)(i).

(1)


- (iii) Give **two** industrial advantages, other than cost, of using ethanoic anhydride rather than ethanoyl chloride in the production of aspirin.

Advantage 1 _____

Advantage 2 _____


(2)

- (d) Complete the following equation for the reaction of one molecule of benzene-1,2-dicarboxylic anhydride (phthalic anhydride) with one molecule of methanol by drawing the structural formula of the single product



(1)

- (e) The indicator phenolphthalein is synthesised by reacting phthalic anhydride with phenol as shown in the following equation.



- (i) Name the functional group ringed in the structure of phenolphthalein.

(1)

- (ii) Deduce the number of peaks in the ¹³C n.m.r. spectrum of phenolphthalein.

(1)

- (iii) One of the carbon atoms in the structure of phenolphthalein shown above is labelled with an asterisk (*).
Use **Table 3** on the Data Sheet to suggest a range of δ values for the peak due to this carbon atom in the ¹³C n.m.r. spectrum of phenolphthalein.

(1)

- (f) Phenolphthalein can be used as an indicator in some acid–alkali titrations.
The pH range for phenolphthalein is 8.3 – 10.0

- (i) For **each** acid.alkali combination in the table below, put a tick (✓) in the box if phenolphthalein could be used as an indicator.

Acid	Alkali	Tick box (✓)
sulfuric acid	sodium hydroxide	
hydrochloric acid	ammonia	
ethanoic acid	potassium hydroxide	
nitric acid	methylamine	

(2)

- (ii) In a titration, nitric acid is added from a burette to a solution of sodium hydroxide containing a few drops of phenolphthalein indicator.

Give the colour **change** at the end-point.

(1)

(Total 21 marks)