


Name of the Student: _____

Max. Marks : 19 Marks

Time : 19 Minutes

Q1.

Iodide ions are oxidised to iodine by hydrogen peroxide in acidic conditions.



The rate equation for this reaction can be written as

$$\text{rate} = k [\text{H}_2\text{O}_2]^a [\text{I}^-]^b [\text{H}^+]^c$$

In an experiment to determine the order with respect to $\text{H}^+(\text{aq})$, a reaction mixture is made containing $\text{H}^+(\text{aq})$ with a concentration of $0.500 \text{ mol dm}^{-3}$

A large excess of both H_2O_2 and I^- is used in this reaction mixture so that the rate equation can be simplified to

$$\text{rate} = k_1 [\text{H}^+]^c$$

- (a) Explain why the use of a large excess of H_2O_2 and I^- means that the rate of reaction at a fixed temperature depends only on the concentration of $\text{H}^+(\text{aq})$.

(2)


- (b) Samples of the reaction mixture are removed at timed intervals and titrated with alkali to determine the concentration of $\text{H}^+(\text{aq})$.

State and explain what must be done to each sample before it is titrated with alkali.

(2)

- (c) A graph of the results is shown in **Figure 1**.

Figure 1



Explain how the graph shows that the order with respect to $\text{H}^+(\text{aq})$ is zero.

(2)

- (d) Use the graph in **Figure 1** to calculate the value of k_1

Give the units of k_1

k_1 _____

Units _____

(3)

- (e) A second reaction mixture is made at the same temperature. The initial concentrations of $\text{H}^+(\text{aq})$ and $\text{I}^-(\text{aq})$ in this mixture are both $0.500 \text{ mol dm}^{-3}$

There is a large excess of H_2O_2


In this reaction mixture, the rate depends only on the concentration of $\text{I}^-(\text{aq})$.

The results are shown in the table.

Time / s	0	100	200	400	600	800	1000	1200
$[H^+] / \text{mol dm}^{-3}$	0.50	0.44	0.39	0.31	0.24	0.19	0.15	0.12

Plot these results on the grid in **Figure 2**. The first three points have been plotted.

Figure 2



(1)

- (f) Draw a line of best fit on the grid in **Figure 2**.


(1)

- (g) Calculate the rate of reaction when $[H^+] = 0.35 \text{ mol dm}^{-3}$
Show your working using a suitable construction on the graph in **Figure 2**.

Rate _____ $\text{mol dm}^{-3} \text{ s}^{-1}$

(2)

- (h) A general equation for a reaction is shown.



In aqueous solution, **A**, **B**, **C** and **D** are all colourless but **E** is dark blue.

A reagent (**X**) is available that reacts rapidly with **E**. This means that, if a small amount of **X** is included in the initial reaction mixture, it will react with any **E** produced until all of the **X** has been used up.

Explain, giving brief experimental details, how you could use a series of experiments to

determine the order of this reaction with respect to **A**. In each experiment you should obtain a measure of the initial rate of reaction.

(6)

(Total 19 marks)