Name of the Student:_____
Max. Marks : 19 Marks **Time : 19 Minutes**

Mark Schemes

Q1.

(a) (i) **M1** drawn curve starts at reactants and ends at products

Tapered lines into the original curve gain credit for M1

M2 curve peak is below the one drawn in the question
(and may show one/two humps)

Mark M1 and M2 independently

2

(ii) Exothermic (reaction)

Ignore "ΔH is negative"

1

(iii) Σ bond (enthalpy) reactants < Σ bond (enthalpy) products

The sum for H_2 and I_2/reactants is less than/lower than/smaller than the sum for 2HI/products
OR
The sum for 2HI/products is more than/larger than/bigger than the sum for H_2 and $I_{2/reactants}$

Accept "It OR the sum will be smaller or less"

1

(iv) **M1** *p*

2

M2 – (*q* – *p*)

OR

p – *q*

OR

– *q* + *p*

M2 demands that the sign for an exothermic reaction is part of the outcome mathematically.
Ignore case

(b) (i) Increase/speed up/faster (rate of attainment of equilibrium)

OR

Increase/speed up/faster rate of both forward and reverse reaction

Downloaded from www.merit-minds.com

OR

Increase/speed up/faster rate of reaction

Credit "It took less time"

1

(ii) **M1** Increase/speed up/faster (rate of attainment of equilibrium)

M2 More particles/molecules in a given volume/space
OR the particles/molecules are closer together
OR an increase in concentration.

M3 More/higher chance of successful/effective/productive collisions (between particles)
OR more collisions/higher chance of collisions (of particles) with $E>E_{Act}$

If M1 is blank, mark on and credit M1 in the text

If M1 is given as "decrease"/"no effect"/"no change" then CE = 0 for clip

In M1, if increase both the forward and reverse reaction, but no mention of rate, penalise M1 but mark on.

In M1, if increase either forward rate or reverse rate only, then penalise M1 but mark on.

Penalise M3 if an increase in the value of E_{Act}/energy of particles is stated.

Max 1 for M2 and M3 if reference to "atoms"

3

[10]

Downloaded from www.merit-minds.com

Q2.

(a) Antacid

OR

to neutralise acidity

OR

eases indigestion

Credit suitable reference to indigestion or to laxative or to relief of constipation

1

(b) **M1** Decrease in T decreases the <u>energy</u> of the <u>particles/ions/H^+/molecules</u>

M2 (also scores M1) Decrease in the number of/less particles/ions/ H^+/molecules with $E \geq E_{Act}$ or $E \geq$ minimum energy to react

*In **M1** and **M2,** credit "atoms" but ignore "calcium carbonate", ignore "calcium", ignore any ion formula except H^+*

M3 Few(er)/Less effective/productive/successful collisions

QoL

3

(c) (i) Strontium has a higher melting point than barium, because

Correct reference to size of cations/proximity of electrons
M1 (For Sr) delocalised <u>electrons closer to cations/positive ions/atoms/nucleus</u>

OR

cations/positive ions/atoms are smaller

OR

cation/positive ion/atom or it has fewer (electron) shells/levels

Ignore general Group 2 statements

Penalise M1 if Sr or Ba is said to have <u>more or less</u> delocalised electrons

Ignore reference to shielding

***CE = 0** for reference to molecules or intermolecular forces or covalent bonds*

Relative strength of metallic bonding
M2 (Sr) has <u>stronger</u> attraction between the <u>cations/positive ions/ atoms/nucleus</u> and the delocalised <u>electrons</u>

OR

<u>stronger metallic</u> bonding
(assume argument refers to Sr but accept converse argument for Ba) 2

Ignore "Van der Waals forces (between atoms)" but penalise if "between molecules"

(ii) $Sr + \mathbf{2}H_2O \rightarrow Sr(OH)_2 + H_2$

Or multiples

Downloaded from www.merit-minds.com

1

(d) **2**Mg + $TiCl_4$ → **2**$MgCl_2$ + Ti

Or multiples

1

[9]

Downloaded from www.merit-minds.com